

---

**Journal of Physical & Theoretical Chemistry**  
**Islamic Azad University of Iran 2(4)**  
(2006)

*Science and Research Campus*

ISSN: 1735-2126

---

**Thermodynamic Analysis for Cationic Surfactants**  
**Binding to Bovine Serum Albumin**

*Abdol-Khalegh Bordbar<sup>a,b1</sup>*

*a: Department of Chemistry, Shahreza Branch, Islamic Azad University, Shahreza, Iran*

*b: Department of Chemistry, Isfahan University, Isfahan, 81746-73441, Iran*

**ABSTRACT**

In the present study, the binding isotherms for interaction of a homologous series of n-alkyltrimethyl ammonium bromides with bovine serum albumin (BSA) have been analyzed on basis of intrinsic thermodynamic quantities. In this regards, the intrinsic Gibbs free energy of binding,  $\Delta G_{b,v}^{(i)}$ , has been estimated at various surfactant concentrations and its trend of variation for both binding sets have been interpreted on basis of cooperativity and hydrophobicity of process. Subsequently, the contribution of electrostatic and hydrophobic interactions in  $\Delta G_{b,v}^{(i)}$ , have been estimated using a published method which has been previously introduced by us for analysis of jack bean urease-cationic surfactant system. The results represent the favoring predominate role of hydrophobic interactions and inhibiting rule of electrostatic interaction in binding affinity of both sets. The predominate role of hydrophobic interactions in the second binding set can be related to entropy statistical effect, which arises from numerous number of binding sites in this set but it may be referred to large amount of positive charge density and accessible hydrophobic surface area of BSA in first binding set.

**Keywords:** Bovine serum albumin; Cationic surfactants; Gibbs free energy of binding; Electrostatic interactions; Hydrophobic interactions

---

<sup>1</sup> Tel: +98-311-7932710 and +98-913-167-7331

Fax+ 98-311-6689732

E-mail: [bordbar@chem.ui.ac.ir](mailto:bordbar@chem.ui.ac.ir) and [khalegh\\_bordbar@yahoo.com](mailto:khalegh_bordbar@yahoo.com)

## 1. INTRODUCTION

In living systems, binding interactions between biopolymers (such as proteins, nucleic acids and poly sachharides) and organic solutes (such as hormones, sugars and fatty acid salts) frequently occur in aqueous media [1]. Such interactions are responsible for the occurrence of many types of bioactive phenomena. Positive binding of many ligands to proteins has been extensively investigated from both the experimental and the theoretical standpoint [2-9]. Excellent reviews on the physicochemical aspects of polymer surfactant interactions have recently been presented by Goddard [10,11]. The different techniques used for such study include equilibrium dialysis, measurement of surface tension, electrical conductivity and viscosity, electrophoresis and ultracentrifugation, gel filtration, ion-specific electrodes, solubilisation, fluorescent probes, electro-optic effects, NMR, small angle neutron scattering, calorimetry, ESR and X-ray diffraction. The effects of surfactant chain length and structure, interaction models and causes for polymer-surfactant complex formation have been discussed in these reviews. It has been suggested that the mechanism of interaction is due to binding charge head groups of the surfactant to the sites with opposite charge at the protein surface, with simultaneous interaction of hydrophobic tail of the surfactant to hydrophobic patches at the protein surface [12]. The above statement of these initial interactions are followed by unfolding and exposure of the hydrophobic interior and hence

generation of numerous hydrophobic binding sites [13,14].

In many studies of protein denaturation and its folding, bovine serum albumin (BSA), which is composed of 583 amino acids and 17 disulfide bonds, has been used with different physicochemical methods [15-18] perhaps, most importantly, due to its well-established primary structure [17, 19]. BSA is largely helical and thermally more stable at pH 7 [18]. Recently, we have investigated the interaction of a series of n-alkyl trimethyl ammonium bromides with BSA using ion selective membrane electrodes as a simple, fast and accurate method [20]. The obtained accurate binding curves have been analyzed on basis of two sets binding sites and the role of both electrostatic and hydrophobic forces have been shown in binding affinity of sites. In the present study, at first, the intrinsic Gibbs free energy of binding has been calculated for both binding sets and its trend of variation has been interpreted on basis of binding mechanism. Subsequently, the contribution of electrostatic and hydrophobic interactions in intrinsic Gibbs free energy has been estimated using an approach which was successfully applied for interaction of cationic surfactants with jack bean urease (JBU), previously [21].

## 2. DATA ANALYSIS AND RESULTS

It has been previously shown that the intrinsic Gibbs free energy of binding per mole of surfactant ions for first  $\Delta G_{b,v}^{(1)}$  and

second,  $\Delta G_{b,v}^{(2)}$  binding set can be calculated from the following formula [22],

$$\Delta G_{b,v}^{(1)} = -RTn_{H1} \ln K_{H1} + RT(1 - n_{H1}) \ln[S]_f$$

if  $0 < v \leq g_1$  (1)

$$\Delta G_{b,v}^{(2)} = -RTn_{H2} \ln K_{H2} + RT(1 - n_{H2}) \ln[S]_f$$

if  $g_1 < v \leq g_1 + g_2$  (2)

$R$ ,  $T$ ,  $[S]_f$  and  $v$  are gas universal constant, absolute temperature, free surfactant concentration and average number of bound surfactant ions per each macromolecule in these formulas, respectively. Where  $g_1$ ,  $n_{H1}$  and  $K_{H1}$  are the number of binding sites, Hill coefficient and the Hill binding constant, for first binding set and  $g_2$ ,  $n_{H2}$  and  $K_{H2}$  are the corresponding parameters for second binding set, respectively. With respect to the nature of interaction,  $\Delta G_{b,v}^{(i)}$  can be considered as a summation of two parts, as follow:

$$\Delta G_{b,v}^{(i)} = \Delta G_{b,v}^{(i)}(ele) + \Delta G_{b,v}^{(i)}(hyd) \quad (3)$$

where  $\Delta G_{b,v}^{(i)}(ele)$  and  $\Delta G_{b,v}^{(i)}(hyd)$  are the electrostatic and hydrophobic contribution to intrinsic Gibbs free energy of binding for  $i$ th set, respectively. For binding of a homologous series of  $n$ -alkyl trimethyl ammonium bromide,  $\Delta G_{b,v}^{(i)}(hyd)$  depends to hydrocarbon tail length of surfactant while  $\Delta G_{b,v}^{(i)}(ele)$  does not. This dependency can be represented by the following relation:

$$\Delta G_{b,v}^{(i)}(hyd) = f(C_n) \quad (3)$$

where  $C_n$  and  $f(C_n)$  are the number of carbon atoms in the hydrocarbon tail of surfactant and any arbitrary function of  $C_n$ , respectively. It is obvious that:

$$\lim_{C_n \rightarrow 0} \Delta G_{b,v}^{(i)}(hyd) = 0 \quad (4)$$

or

$$\lim_{C_n \rightarrow 0} \Delta G_{b,v}^{(i)} = \Delta G_{b,v}^{(i)}(ele) \quad (5)$$

This simple idea can be used for estimation of  $\Delta G_{b,v}^{(i)}(ele)$  and  $\Delta G_{b,v}^{(i)}(hyd)$ . Figs. 1 and 2 show the variation of  $\Delta G_{b,v}^{(1)}$  and  $\Delta G_{b,v}^{(2)}$  versus  $\log[S]_f$  for interaction of dodecyl trimethyl ammonium bromide (DTAB), tetradecyl ammonium bromide (TTAB) and hexadecyl trimethyl ammonium bromide (HTAB) with BSA, respectively. The required data for calculation of  $\Delta G_{b,v}^{(i)}$  have been directly taken from previous study [20]. The values of  $\Delta G_{b,v}^{(i)}$  at any specified value of  $[S]_f$  have been extracted from these figures and plotted versus  $C_n$  (Figs. 3 and 4). The points relate to the specified value of  $[S]_f$  were fitted in a linear equation using least-square fitting program. With respect to Eq. (5),  $\Delta G_{b,v}^{(i)}(ele)$  should be equal to Y-intercept of these lines, and subsequently, the values of  $\Delta G_{b,v}^{(i)}(hyd)$  can be estimated by subtracting of  $\Delta G_{b,v}^{(i)}(ele)$  from  $\Delta G_{b,v}^{(i)}$ . Figs. 5 and 6 show the variation of

$\Delta G_{b,v}^{(i)}(ele)$  and  $\Delta G_{b,v}^{(i)}(hyd)$  versus  $\log[S]_f$  respectively.

for first and second binding sets,




Fig. 1. The variation of  $\Delta G_{b,v}^{(1)} (kJ/mol)$  vs.  $\log[S]_f$  for interaction of BSA with DTAB ( $\circ$ ), TTAB ( $\bullet$ ) and HTAB ( $\blacktriangle$ ).






Fig. 2. The variation of  $\Delta G_{b,v}^{(2)} (kJ/mol)$  vs.  $\log[S]_f$  for interaction of BSA with DTAB ( $\circ$ ), TTAB ( $\bullet$ ) and HTAB ( $\blacktriangle$ ).



**Fig. 3.** The variation of  $\Delta G_{b,v}^{(1)} (kJ/mol)$  vs.  $C_n$  for interaction of BSA with cationic surfactants at various  $[S]_f$ . (a)  $9.98 \times 10^{-5}$  M, (b)  $12.57 \times 10^{-5}$  M, (c)  $15.82 \times 10^{-5}$  M, (d)  $19.92 \times 10^{-5}$  M and (e)  $25.08 \times 10^{-5}$  M of surfactant (the arrow shows the direction of surfactant concentration increasing).



**Fig. 4.** The variation of  $\Delta G_{b,v}^{(2)} (kJ/mol)$  vs.  $C_n$  for interaction of BSA with cationic surfactants at various  $[S]_f$ . (a)  $2.34 \times 10^{-3}$  M, (b)  $1.73 \times 10^{-3}$  M, (c)  $2.00 \times 10^{-3}$  M, (d)  $2.24 \times 10^{-3}$  M and (e)  $2.51 \times 10^{-3}$  M of surfactant (the arrow shows the direction of surfactant concentration increasing).




Fig. 5. The variation of  $\Delta G_{b,v}^{(1)}(ele)(kJ/mol)$  ( $\square$ ) and  $\Delta G_{b,v}^{(1)}(hyd)(kJ/mol)$  for interaction of BSA with DTAB ( $\circ$ ), TTAB ( $\bullet$ ) and HTAB ( $\blacktriangle$ ) with  $\log[S]_f$ .




Fig. 6. The variation of  $\Delta G_{b,v}^{(2)}(ele)(kJ/mol)$  ( $\square$ ) and  $\Delta G_{b,v}^{(2)}(hyd)(kJ/mol)$  for interaction of BSA with DTAB ( $\circ$ ), TTAB ( $\bullet$ ) and HTAB ( $\blacktriangle$ ) with  $\log[S]_f$ .

### 3. Discussion and conclusion

The negative slope of the lines in Figs. 1 and 2 represent the positive cooperativity in the binding process. The positive cooperativity in both sets can be related to

special role of hydrophobic forces in the formation of BSA-surfactant complexes. However, the more steepness of the lines in Fig. 1 represents the more predominate rule of hydrophobic interactions in the first binding set. This observation is in

contradiction with JBU – cationic surfactants system. With respect to these observations, the following assumptions of the model regarding the changes in the state of the protein at different concentrations of surfactant can be defined as follows: the first type of binding sites is present in the native protein. The saturation of these binding sites is cooperative, then, surfactant binding induce a considerable change in conformational state of BSA. This large conformational change can be occurred with unfolding and exposure of numerous non-specific binding sites.

The more negative values of  $\Delta G_{b,v}^{(1)}$  with respect to  $\Delta G_{b,v}^{(2)}$  for any surfactant, represents the stronger initial interactions, which are usually due to attractive electrostatic forces between cationic head group of surfactant ions with negative charge centers at the protein surface. The most interesting part of this paper is the use of experimental data (obtained with a series of ionic surfactants differing in the length of the hydrocarbon tail) to obtain the hydrophobic and electrostatic components of binding energies. In this regard the novel feature of Figs. 4 is the existence of iso-affinity point. This point is at  $C_n$  equal to 19.57. It seems that this point is inflection point for kind of cooperativity. The meaning of isoaffinity point in Fig. 4 can be interpreted as follows: It is well known that denaturation power of ionic surfactant increased by increasing of hydrocarbon tail. However, the stability of protein has a limited value so that it is expected that after a specified value of  $C_n$  all of the surfactants with various tail length behave identically. Hence, it can be suggested that these two limiting values for  $C_n$  of surfactant relate to the denaturing power of homologous surfactants and the extent of structural stability of protein. However, such inflection point has not been observed in Fig.3 that corresponds to first binding set.

This is in contradiction with JBU-surfactant system. Fig. 5 represents that the contribution of electrostatic interactions is less than hydrophobic in the first binding set for all of the surfactants. Moreover, an inhibition effect is observed for electrostatic interactions. So, the predominant driving force in first and second binding sets is hydrophobic interactions. However, the variation trend of  $\Delta G_{b,v}^{(1)}(ele)$  to less positive values is not in agreement with increasing of positive charge density in the BSA due to binding by cationic surfactants. This may be related to conformational changes of BSA that reduces the positive charge density on BSA. The values of  $\Delta G_{b,v}^{(1)}(hyd)$  are going up to more negative values due to increasing of  $C_n$  or hydrocarbon tail length, which is expected.

With respect to Fig. 6 the positive values of  $\Delta G_{b,v}^{(i)}(ele)$  represents the net positive charge in protein in all binding stages of second binding set. In the other word, the repulsive electrostatic forces between cationic head group of surfactant and positive charges in the BSA-surfactant complexes inhibited the binding of next surfactant ions.

However, these values are less positive corresponds to first binding set. This may be related to unfolding of protein and reduces of positive charge density on protein. The values of  $\Delta G_{b,v}^{(i)}(hyd)$  are negative and represents the favoring effects of hydrophobic interaction in binding process for both binding sets and its values is sufficient that can compensate the repulsive electrostatic forces, effectively. It can be concluded that hydrophobic interactions have an essential role in binding process of cationic surfactant to BSA. Part of this role can also be related to the numerous numbers of binding sites in the second binding set, which increased the statistical entropy part of macroscopic

Gibbs free energy. However, in comparison with JBU-surfactant, it looks that the role of hydrophobic interactions is much more, especially in first binding set.

#### Acknowledgements

The financial support of Research Council of Shahreza-Branch of Islamic Azad University is gratefully acknowledged.

#### References

- [1] Stryer, L., Biochemistry, 4th edn., W.H. Freeman, New York, 1994.
- [2] Steinhardt, J. and Reynolds, J.A., Multiple equilibria in proteins, Academic Press, New York, 1969.
- [3] Steinhardt, J., Stocker, N. and Birdi, K.S., Biochemistry, 13 (1974) 4461.
- [4] Reboiras, M.D., Pfisher, H. and Pauly, H., Biophys. Chem., 9 (1978) 37.
- [5] Manning, G.S., Biophys. Chem., 9 (1978) 65.
- [6] Gilanyi, T. and Wolfram, E., Colloids Surf., 3 (1981) 181.
- [7] McGhee, J.D. and von Hippel, P.H., J. Mol. Biol., 86 (1974) 469.
- [8] Gill, S.J., Gaud, H.T., Wyman, J. and Barias, B.G., Biophys. Chem., 8 (1978) 53.
- [9] Spector, A.A. and Santos, E.C., Ann. N.Y. Acad. Sci., 226 (1973) 247.
- [10] Goddard, E.D., Colloids Surf., 19 (1986) 301.
- [11] Goddard, E.D. and Mittal, K.L., Surfactants in solution, Plenum Press, New York, 1992.
- [12] Jones, M.N., Manley, P., Chem. Soc. Faraday Trans. 1 76 (1980) 654.
- [13] Tipping, E., Jones, M.N. and Skinner, H.A., Chem. Soc. Faraday Trans. 1 70 (1974) 1306.
- [14] Oakes, J., Chem. Soc. Faraday Trans. 170 (1974) 2200.
- [15] Oakes J. (1973) Magnetic resonance studies of the interaction between bovine serum albumin and surfactants. Eur. J. Biochem. 36, 553--558.
- [16] Jones M. N., Skinner H. A. and Tipping E. (1975) The interaction between BSA and surfactants. Biochem. J. 47, 229 - 234.
- [17] Batra P. P., Sasa K. T. and Takeda K. (1989) Circular dichroic study of the conformational stability of sulfhydryl blocked BSA. Int. J. Biochem. 21, 857-862.
- [18] Yamasaki M., Yana H. and Aoki K. (1990) DSC studies on BSA: Effects of pH and ionic strength. Int. J. Biol. Macromolec. 12, 263-268.
- [19] Hilak M. C., Harmsen B. J. M., Braam W. G. M., Joordens J. M. and van OS G. A. J. (1974) Conformational studies on large fragments of bovine serum albumin in relation to the structure of the molecule. Int. J. Pept. Prot. Res. 6, 955-101.
- [20] Rafati, A. A., Bordbar, A. K., Gharibi, H., Amini, M. K. and Safarpour, M. A, Bull. Chem. Soc. Jpn., 2004,
- [21] Bordbar, A. K., Sohrabi, N. and Hojjati, E., Colloids and Surfaces B: Biointerfaces 39 (2004) 171-175
- [22] Bordbar, A. K., Saboury, A. A., Housaindokht, M. R., Moosavi-Movahedi, A. A., J. Colloid Interface Sci. 192 (1997) 415-419.